

Genetic Characterization Based on RAPD-PCR in Cultured Strains of *Clarias gariepinus* (Siluriformes: Clariidae)

Ola-Oladimeji Folasade Adesola^{1,*} , Michael Olufemi Awodiran², Victor Folorunso Olaleye², Joel Idowu Awopetu²

¹Ekiti State University, Department of Zoology and Environmental Biology, P.M.B. 5363, Ado-Ekiti, Ekiti State, Nigeria.
²Obafemi Awolowo University, Department of Zoology, P.M.B. 13, Ile-Ife, Osun State, Nigeria.

Article History

Received 30 April 2020
Accepted 26 August 2020
First Online 07 September 2020

Corresponding Author

Tel.: +2348034797058
E-mail:
folasade.majolagbe@eksu.edu.ng

Keywords

Polymerase
Variability
Population
Dendrogram
Primers

Abstract

The genetic polymorphisms of catfish populations collected from four fish farms in four Southwestern states of Nigeria were determined by Randomly Amplified Polymorphic DNA Polymerase Chain Reaction technique. This was to characterize the strains genetically and provide information on the studied catfish. Following standard procedures, polymerase chain reaction was performed by mixing 15 µl reaction mixture containing FIREPOL[®] Taq DNA polymerase, MgCl₂, dNTPs, distilled water, dyes and glycerol, with 1.0 µl DNA sample of 16-week old *C. gariepinus* strains using OPA03, OPA04, OPC02, OPB08, OPC11, OPG16, OPG19 and OPA19 primers. A sum of 1708 loci having 3072 bands was amplified in all the samples. The RAPD analysis revealed significant genetic variability ($P < 0.05$) among the four sampled populations. The unweighted pair group method with average (UPGMA) dendrogram based on Nei's unbiased genetic distance matrix separated the *Clarias gariepinus* populations into two clades. The first clade was made up of populations from Ibadan and Abeokuta while the second clade consisted of populations from Ado-Ekiti and Ile-Ife. Thus, it is imperative to determine the genetic variation and population structure of the stocks of *C. gariepinus* in advance before commencing on breeding programmes.

Introduction

Clarias gariepinus is a desired aquaculture fish in Nigeria. It belongs to the Kingdom Animalia, Phylum Chordata, Class Actynopteri, Order Siluriformes and Family Clariidae (Freyhof, 2014). It is a member of the most diverse genus, *Clarias* (Legendre, Teugels, Cauty, & Jalabert, 1992). It has a helmet-like head and scaleless, bony elongated body with long dorsal and anal fins. It is a typical air-breathing catfish that has a broad terminal mouth with four pairs of barbels. Its colour varies dorsally from dark to light brown and is often speckled with shades of olive and grey while the lower side is pale cream to white (Skelton, 2001).

Since this fish is a widely accepted species, scientific studies on it cannot be over-emphasised. Even though traditional methods like morphological (Ola-

Oladimeji, et al., 2017; Okomoda, Koh, Hassan, Amornsakun, & Shahreza, 2018) and biochemical studies ((Ola-Oladimeji, Idowu, Adewumi, & Fafowora, 2018) which have been reported to be laborious and time-consuming (Coulo, Vessen, Hofstra, & Huisveld, 1994) and are less efficient in detecting differences in traits have been carried out on the fish, some studies have also been done on its genetic management like hybridization (Ola-Oladimeji, 2015) and cytogenetic techniques such as polyploidy, androgenesis and gynogenesis (Dunham, 2011; Olaniyi & Omitogun, 2012; Karami et al., 2016; Okomoda et al., 2020).

Recently, the application of molecular primers based on the relative difference in deoxyribonucleic acid (DNA) sequences between individuals has been shown to detect more polymorphisms than morphological and protein-based markers. Molecular primers are also used

to study genomes, isolate genes, locate genes on chromosomes, determine gene expression, conduct gene-linkage maps, study the molecular mechanisms of performance, conduct population genetics analysis and apply marker-assisted selection (Liu & Dunham, 1998).

It is well-known that all inherited characteristics or traits are under genetic control; consequently, various methods have been developed for genetic analysis of populations. The molecular analyses are the most direct and potent, and these were enhanced by the development of the Polymerase Chain Reaction (PCR). The PCR depends on the ability to alternately denature (melt) double-stranded DNA molecules and renature (anneal) complementary single strands in a controlled fashion (Lodish et al., 2016). The variations in the size of a gene can then be detected by a change in the size of the PCR product which can be determined by using gel electrophoresis (Dale & Park, 2004).

According to Ikpeme et al. (2015), genetic analysis can be used to determine variation. Danish & Singh (2018) also found that significant population differentiation which indicated some degree of intra- and inter-population genetic variations existed in two populations of *Clarias batrachus*. This was suggested to be due to the difference in habitat and breeding strategies between the two populations. One of several markers used in the analyses is Random Amplified Polymorphic DNA (RAPD). This technique uses short primers of arbitrary nucleotide sequence to access and reproducibly amplify random segments of genomic DNA (from a wide variety of species) to reveal polymorphisms among the amplification products which are useful as genetic markers and can be detected through

examination of an ethidium bromide-stained agarose gel (Williams, Kubelik, Livak, Rafalski, & Tingey, 1990). The RAPD markers are for various uses in the detection of genetic diversity in population genetics (Zheng et al., 2019) and it was used to characterize cultured strains of *C. gariepinus* in this study.

A few studies (Saad, Hanafi, Essa, Guerges, & Ali, 2009; Popoola, Fasakin, & Awopetu, 2014; Ikpeme et al., 2015) have been reported on genetic similarity and diversity of different populations of *C. gariepinus*. In spite of these, information on the genetic characterization of the fast- and slow-growing strains of cultured *C. gariepinus* using Randomly Amplified Polymorphic DNA is yet to be provided. Therefore, it is important to study these strains of catfish more because fish farmers are faced with challenges of managing runts during culture and these have reduced their productivity and income.

This study aims to characterize the cultured strains of *C. gariepinus* in Southwestern Nigeria genetically using Random Amplified Polymorphic DNA (RAPD) markers for management and preservation of its genetic resources in future studies.

Materials and Methods

Identification and Collection of Fish Specimens

Forty (40) eight-week-old fast- and 40 slow-growing strains of *C. gariepinus* were identified, collected from four selected farms in four Southwestern states of Nigeria (Figure 1) for comparison and to obtain a better result. The two strains were obtained by selecting and




Figure 1: Map of Nigeria with colour-coded states where the *C. gariepinus* juveniles were collected.

separating the shooters from the runts after two weeks of rearing. Samples were collected from the following selected locations; the Teaching and Research Farm, Federal University of Agriculture, Abeokuta (FUNAAB) (7° 09' 20" N & 3° 20' 42" E); Sun Fish Farm, Ibadan, (7° 22' 39" N & 3° 54' 21" E); Ade Aquaculture Centre, Ado-Ekiti, (7° 37' 23" N & 5° 13' 15" E); and Edens Fish Farm, Ile-Ife, (7° 28' 0" N & 4° 34' 0" E). At the sampling sites, each of the fast- and slow-growing strains was separated appropriately and the fish were raised in different tanks simultaneously at Edens Fish Farm, Ile-Ife.

Blood Sample Collection for DNA Extraction

Genomic DNA was extracted from the blood of each of the individual fish (40 in total) from the selected groups after rearing using NORGEN Blood Genomic DNA Isolation Kit (NORGEN, Biotechnology Corporation, Canada, 2014). The extraction was performed according to El-Mogy, Simkin, and Haj-Ahmad (2016), using spin column chromatography method.

RAPD and Electrophoresis

DNA amplification was done using RAPD (OPA03, OPA04, OPC02, OPB08, OPC11, OPG16, OPG19 and OPA19) according to standard but a modified method of Williams et al. (1990). Table 1 shows the characterization of the primers used including the primer sequences and annealing temperatures. The amplified products (8 µl) of each sample were separated by electrophoresis in 1% agarose gels buffered with 1X TAE and the polymorphisms among the amplification products were detected by staining with 15 µl of 0.5 µg/mL Ethidium Bromide. A PCRSizer 100 bp DNA ladder (100bp - 1000bp) (Norgen Biotek Corporation, Canada) was electrophoresed alongside the RAPD reactions as molecular weight marker. Electrophoresis was run at 100 V for 2.5 hours. Finally, the DNA bands were examined under Ultra-Violet (UV) transilluminator system (ZENITH, India), and the stained gels were photographed and saved on the computer for further analysis.

Statistical Analysis

The results obtained after rearing the catfish were analysed with Descriptive statistics, Analysis of variance

(ANOVA) and Duncan's multiple range test (DMRT) using IBM SPSS Statistics (Version 22, 2013) software.

Scoring of RAPD Bands

The RAPD bands were scored from the photomicrographs of the gel as binary data using GelQuest-DNA Fragment Analysis Software, Version 3.2.1 (©2010 SequentiX-Digital DNA Processing, Germany). A single data matrix was constructed pooling the binary data (1/0) for all the primers.

Data Analyses for RAPD

GenAlEx 6.502 genetic software (Peakall & Smouse, 2006; 2012) was used to analyse the data generated to compute the number of different bands, analysis of molecular variance (AMOVA), Nei's genetic identity and Nei's genetic distances. Nei's genetic distances were calculated for all population pairs to construct a phylogenetic tree (Unweighted pair group method with average dendrogram) using MEGA evolutionary genetic analysis software (Version 7.0.21, 2018).

Results

Length and Weight Traits of *C. gariepinus*

The mean total length and body weight of the two strains of *C. gariepinus* used for DNA analysis are shown in (Table 2). The mean total length and body weight were significantly higher at $P < 0.05$ in the fast-growers than in the slow-growing strains.

Profile of RAPD Markers

Plate 1 shows the representative agarose gel electrophoretic profile of RAPD using OPG16 primer. For all the primers used, a sum of 1708 loci having 3072 bands was amplified in all the samples. The fast-growing catfish had 1522 bands while slow-growing fish showed 1550 bands (Table 3). In Table 4, the lowest number of private bands (85) among the slow-growing population was recorded in the Ibadan population, while among the fast-growers; the population from Abeokuta had the least (74). The analysis showed that the number of locally common bands were correspondingly higher in

Table 1. Characterization of RAPD primers used in the analysis, including primer sequences and annealing temperatures (Ta (°C))

Primers	Sequences	Ta (°C)
OPA03	5'AGTCAGCCAC3'	37.0
OPA04	5'AATCGGGCTG3'	35.1
OPC02	5'GTGAGGCGTC3'	37.6
OPB08	5'GTCCACACGG3'	37.3
OPC11	5'AAAGCTGCGG3'	36.9
OPG16	5'AGCGTCTCC3'	38.8
OPG19	5'GTCAGGGCAA3'	34.7
OPA19	5'CAAAGTCGG3'	32.2

Table 2. Differences in length and weight between the two strains of *C. gariepinus* used for DNA analysis

Strains	Total length (cm)	Body weight (g)
Fast-growing <i>C. gariepinus</i>	40.44 ^a ± 0.58	466.66 ^a ± 18.66
Slow-growing <i>C. gariepinus</i>	21.55 ^b ± 0.50	81.89 ^b ± 5.62

*Means in the same column superscripted by different letters differed significantly (P<0.05)

Table 3. Number of bands and loci

Population type	Primers								Total
Slow-growers									
	OPA03	OPA04	OPA19	OPB08	OPC02	OPC11	OPG16	OPG19	
SAB	34	36	31	36	37	41	52	40	307
SIB	27	32	35	45	29	48	42	38	296
SAD	114	46	125	36	43	45	42	5	456
SIL	126	49	137	38	37	36	44	24	491
									1550
Fast-growers									
FAB	32	30	34	46	47	0	43	33	265
FIB	26	34	34	45	30	46	47	40	302
FAD	128	42	133	42	37	49	46	0	477
FIL	106	46	125	45	41	49	47	19	478
Total									1522
Total Number of loci	286	188	304	193	181	205	213	138	1708

Table 4. Band patterns obtained after band scoring across the populations

Population	No. private bands	No. shared bands (<=25%)
Slow-growers		
SAB	94	102
SIB	85	119
SAD	123	139
SIL	132	154
Fast-growers		
FAB	74	92
FIB	96	92
FAD	111	164
FIL	136	150

No. private bands = No. of bands unique to a single population

No. shared bands (<=25%) = No. of common bands (freq. >= 5%) found in 25% or fewer populations

SAB= Slow-growers Abeokuta, **SIB**= Slow-growers Ibadan, **SAD**= Slow-growers Ado-Ekiti, **SIL**= Slow-growers Ile-Ife. **FAB**=Fast-growers Abeokuta, **FIB**= Fast-growers Ibadan, **FAD**= Fast-growers Ado-Ekiti, **FIL**= Fast-growers Ile-Ife

the slow-growing populations from across locations than those of the fast-growers. The only exception recorded was in the Ado-Ekiti population where the slow-growers had lower (139) locally common bands than the fast-growers (164).

Plate 1(A and B) Representative agarose gel electrophoretic profile of RAPD using OPG16 primer for adult *C. gariepinus* (Figure 2)

Population Differentiation and Structure

The Analysis of Molecular Variance, AMOVA showed the significance levels of genetic variability among regions, Φ_{PR} and among regions corrected for

bias Φ'_{PR} were 0.01. Most of the genetic diversity occurred within populations (92%) while the variability among regions and populations contributed 7% and 1% respectively (Figure 3, Table 5).

The value of inter-population Nei's genetic identity was found least (0.963) between slow-growers from Ibadan and fast-growers from Ado-Ekiti. The highest (0.985) value was recorded between the strains from Abeokuta (Table 6). Also, the unweighted pair group method with average (UPGMA) dendrogram based on Nei's unbiased genetic distance matrix (Nei, 1972) performed according to RAPD analysis separated the four catfish populations sampled into two clades. The first clade was formed by populations sampled from

Ibadan and Abeokuta while the second was formed by populations sampled from Ado-Ekiti and Ile-Ife (Figure 4).

Discussion

This study has shown the existence of genetic variability and relatedness among and within *C. gariepinus* populations from four hatcheries in South-Western Nigeria. The catfish have shown some similarities with regards to the number of bands shared in common and variations have also been observed as a result of some fragments present in one group and absent in another in the locally common bands.

According to Awodiran and Afolabi (2018), genetic variations, which are known as 'the raw materials for evolution,' are of importance in the survival of species. There is a need to deliberately introduce new breeds

into the cultured populations in order to enhance the evolutionary potentials of these populations. To avoid a vicious cycle, the introduction of these new genotypes must be done methodically, because outbreeding depression is an impending risk if the genetic distance between the populations and the new breeds is too large.

The high genetic identity observed in this research may be because these catfish are from cultured environments. These were comparable with Popoola et al. (2014) and Ikpeme et al. (2015) who worked on *C. gariepinus*. While Ikpeme et al. (2015) reported that as genetic distance increased, genetic identity narrowed between the wild and the cultured species, Popoola et al. (2014) reported that the percentage of genetic identity and genetic distance of RAPD-PCR profile among six populations of wild and cultured *C. gariepinus* ranged from 74.6% to 83.5% and from 0.180 to 0.293




Figure 2. Plate 1(A and B): Representative agarose gel electrophoretic profile of RAPD using OPG16 primer for adult *C. gariepinus*.

Lanes 1-5= FAB, 6-10= SAB, L = DNA ladder, 11-15=FIB, 16-20=SIB. Lanes 21-25= FAD and 26-30= SAD, L = DNA ladder, lanes 31-35= FIL and 36-40=SIL.

FAB= Abeokuta fast-growers, SAB= Abeokuta slow-growers, FIB=Ibadan fast-growers, SIB=Ibadan slow-growers, FAD= Ado-Ekiti fast-growers, SAD= Ado-Ekiti slow-growers, FIL= Ile-Ife fast-growers, SIL= Ile-Ife slow-growers.

Table 5. Analysis of molecular variance (AMOVA) showing the partitioning of genetic variation among regions and populations, and within populations of *C. gariepinus* studied

Source of variation	Est. var.	%	Φ PR	Φ' PR	P(rand >= data)
Among Regions	5.878	7%			
Among Populations	1.097	1%			
Within Population	83.938	92%			
Total	90.913	100%	0.01	0.01	0.0001

Est. var. = Estimate of variance

%= Percentage of total variation

Φ PR = Genetic variability among regions

Φ' PR = Genetic variability among regions corrected for bias

P(rand >= data) = Level of significance




Figure 3. Sources and proportion of molecular variance: Input as binary distance matrix.

Table 6. Pairwise population matrix of Nei's genetic identity obtained after analysing RAPD bands

FAB	SAB	FIB	SIB	FAD	SAD	FIL	SIL	
1.000								FAB
0.985	1.000							SAB
0.984	0.984	1.000						FIB
0.983	0.983	0.982	1.000					SIB
0.966	0.964	0.966	0.963	1.000				FAD
0.970	0.969	0.969	0.968	0.971	1.000			SAD
0.969	0.968	0.968	0.967	0.967	0.971	1.000		FIL
0.968	0.966	0.967	0.965	0.966	0.970	0.971	1.000	SIL




Figure 4. Genetic relationship among different populations of *C. gariepinus* using RAPD analysis.

respectively (Popoola, et al., 2014). In this study, the levels of significance of genetic variability among regions ($\Phi_{PR} = 0.01$) indicated that the four sampled populations were significantly different from each other at $P=0.01$, but not at higher probability levels. Pair wise population matrix of Nei's genetic distance obtained after analysing RAPD bands gave the extent of gene differences among populations. The index was used to construct a dendrogram which summarized evolutionary

relationships among the populations into two clades. Also, a number of studies on genetic relationship had been done on fish, one of these include Ogbuebunu & Awodiran (2017) on the genetic relationship among three populations of *Lates niloticus* depicted by UPGMA dendrogram. In the present study, the two clades formed by the UPGMA dendrogram showed that populations sampled from Ibadan and Abeokuta were more closely related than those obtained from Ado-Ekiti

and Ile-Ife. Also, the two different strains obtained from the same location were more related than the others. This relation follows geographical distance. A similar report had been made by Saad et al. (2009) on some of the populations of *C. gariepinus* obtained from the wild.

Analysis of Molecular Variance (AMOVA) has been commonly employed for hierarchical analysis of the genetic differentiation among populations (Wollebaek, Heggenes & Roed, 2011). In this study, the levels of significance of genetic variability among regions indicated that the four sampled fast- and slow-growing populations of adult *C. gariepinus* were significantly different from each other. This is similar to the report of Awodiran and Afolabi (2018) on the AMOVA for RAPD marker analysed on wild and cultured populations of *C. gariepinus* which indicated significant difference ($P=0.01$) between the two populations. Evaluation of the distribution of molecular variation by AMOVA which also illustrated that most of the total genetic variance was found within strains was similar to Ribolli, Rodrigues de Melo and Zaniboni-Filho (2012) who reported that several individuals belonging to a particular population were attributed to other populations.

In conclusion, it was validated in this study that RAPD markers can evaluate genetic variability within cultured populations of *C. gariepinus*. This has corroborated the existing information regarding the need for determining genetic variation and population structure of the stocks of *C. gariepinus* in advance before commencing on breeding programmes. However, markers such as microsatellite DNA can be assayed in further studies for better results.

Acknowledgment

The authors thanked Afolabi O. for his assistance on the DNA analyses.

References

- Awodiran, M.O. & Afolabi, O. (2018). Genetic diversity in cultured and wild population of *Clarias gariepinus* (Burchell, 1822) in Nigeria Using Random Amplified Polymorphic DNA (RAPD) and Microsatellite DNA. *Fisheries and Aquaculture Journal*, 9: 247. <https://doi.org/10.4172/2150-3508.1000247>
- Coulo, M.M.B., Vessen, J.M.B.M., Hofstra, H., & Huisveld, J.H.J. (1994). RAPD analysis: a rapid technique for differentiation of spoilage yeasts. *International Journal of Food Microbiology*, 24(1): 249-260
- Dale, J.W., & Park, S.F. (2004). *Molecular Genetics of Bacteria* (4th Edition). John Wiley and Sons Ltd, England. 346 pp.
- Danish, M. & Singh, I.J. (2018). Assessment of genetic diversity of two populations of catfish *Clarias batrachus* L. using random amplified polymorphic DNA (RAPD) markers. *Journal of Pharmacognosy and Phytochemistry*, 7(3): 2345-2352. <http://dx.doi.org/10.22271/phyto>
- Dunham, R.A. (2011). *Aquaculture and Fisheries Biotechnology: Genetic Approaches*. Second Edition. CAB International, Wallingford, UK. 495 pp.
- El-Mogy, M., Simkin, M., & Haj-Ahmad Y. (2016): Comparative study of DNA isolated from increasing volumes of blood using Norgen's blood genomic DNA isolation kit versus Qiagen's QIAamp DNA blood mini kit. *Norgen Biotechnology Corporation: Application Note 59 Blood DNA Preparation*. Pp. 1-3.
- Freyhof, J. (2014). *Clarias gariepinus*. *The IUCN Red List of Threatened Species* 2014: e.T166023A1106307. <https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T166023A1106307.en>. Downloaded on 12 June 2020.
- Ikpeme, E.V., Udensi, O.U., Ekaluo, U.B., Kooffreh, M.E., Okolo, C.M., Ekpo, P.B. & Ogbonna, N.C. (2015). Unveiling the genetic diversity in *Clarias gariepinus* (Burchell 1822) using Random Amplified Polymorphic DNA (RAPD) fingerprinting technique. *Asian Journal of Animal Sciences*, 9: 187-197. <https://doi.org/10.3923/ajas.2015.187.197>
- Karami, A., Goh, Y.M., Jahromi, M.F., Lazorchak, J.M., Abdullah, M., & Courtenay, S.C. (2016). Diploid and triploid African catfish (*Clarias gariepinus*) differ in biomarker responses to the pesticide chlorpyrifos. *The Science of the Total Environment*, 557-558:204-211. <https://doi.org/10.1016/j.scitotenv.2016.03.030>
- Legendre, M., Teugels, G.G., Cauty, C. & Jalabert, B. (1992). A comparative study on morphology, growth rate and reproduction of *Clarias gariepinus* (Burchell 1822), *Heterobranchus longifilis* Valenciennes, 1840 and their reciprocal hybrids (Pisces, Clariidae). *Journal of Fish Biology*, 40:59-79. <https://doi.org/10.1111/j.1095-8649.1992.tb02554.x>
- Liu, Z.J., & Dunham, R. (1998). Genetic Linkage and QTL Mapping of Ictalurid Catfish. *Circular Bulletin* 321. Alabama Agricultural Experiment Station, Auburn University, Auburn, Alabama. 19 pp.
- Lodish H., Berk A., Kaiser, C.A., Krieger M., Bretscher A., Ploegh H., Martin, K.C. (2016). *Molecular Cell Biology*. 8th Edition, W.H. Freeman and Company, USA. Pp. 1170.
- Ogbuebunu, K.E., & Awodiran, M.O. (2017). Molecular Characterization of *Lates niloticus* (Perciformes, Latidae) Populations from Tree Nigerian Waterbodies using Random Amplified Polymorphic DNA and Microsatellite Markers. *Vestnik zoologii*, 51(1): 31-36. <https://doi.org/10.1515/vzoo-2017-0005>
- Okomoda, V.T., Aminath L., Oladimeji S.A., Abol-Munaf, A.B., Korede, A.I., Ikhwanuddin Mhd, Shahreza, S.M. (2020). First Report on Successful Triploidy Induction in *Clarias gariepinus* (Burchell, 1822) Using Electroporation. *Scientific Reports*, 10: 2425. <https://doi.org/10.1038/s41598-020-59389-2>
- Okomoda, T.V., Koh, I.C.C., Hassan, A., Amornsakun, T., & Shahreza, S.M. (2018). Morphological characterization of the progenies of pure and reciprocal crosses of *Pangasianodon hypophthalmus* (Sauvage, 1878) and *Clarias gariepinus* (Burchell, 1822). *Scientific Reports*, 8(1):3827. <https://doi.org/10.1038/s41598-018-22149-4>
- Ola-Oladimeji F.A. (2015). Comparison of Fitness Traits between Reciprocal Hybrids and Parentals of *Clarias gariepinus* (Burchell, 1822) and *Heterobranchus bidorsalis* (Geoffroy Saint-Hilaire, 1809). *IOSR Journal of Pharmacy and Biological Sciences*, 10(5): 78-82. <https://doi.org/10.9790/3008-10527882>
- Ola-Oladimeji, F.A., Idowu, E.O., Adewumi, A.A., & Fafowora, K.C. (2018). Heterogeneity Studies of Wild *Clarias gariepinus* (Osteichthyes, Clariidae) Using SDS-Polyacrylamide Gel Electrophoresis. *Vestnik Zoologii*, 52,

- 457 - 462. <https://doi:10.2478/vzoo-2018-0047>
- Ola-Oladimeji, F.A., Oso, J.A., Oladimeji, T.E., Idowu, E.O., Adeleke, K. & Urihe, F.O. (2017). Phenotypic diversities of four populations of *Clarias gariepinus* (Siluriformes, Clariidae) obtained from Ogun and Ondo State waterbodies in Southwestern Nigeria. *Vestnik zoologii*, 51(4): 285–294. <https://doi:10.1515/vzoo-2017-0034>.
- Olaniyi, W.A. & Omitogun, O.G. (2012). Induction of diploid gynogoneic larvae of African catfish, *Clarias gariepinus* Burchell 1822. *Ife Journal of Agriculture*, 25:73–82.
- Peakall, R. and Smouse, P.E. (2006). GenALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, 6(1): 288-295. <https://doi.org/10.1111/j.1471-8286.2005.01155.x>
- Peakall, R. and Smouse, P.E. (2012). GenALEX 6.5: Genetic analysis in Excel. Population genetic software for teaching and research- an update. *Bioinformatics*, 28: 2537-2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Popoola, O.M., Fasakin, E.A., & Awopetu, J.I. (2014). Genetic variability in cultured and wild populations of *Clarias gariepinus* (Osteichthys: Clariidae) using Random Amplified Polymorphic DNA (RAPD) marker. *Croatian Journal of Fisheries*, 72(1): 5-11.
- Ribolli J., Rodrigues de Melo, C.M. and Zaniboni-Filho, E. (2012). Genetic characterization of the Neotropical catfish *Pimelodus maculatus* (Pimelodidae, Siluriformes) in the Upper Uruguay River. *Genetics and Molecular Biology*, 35 (4): 761-769 <https://doi:10.1590/S1415-47572012005000060>
- Saad, Y.M., Hanafi, M.S., Essa, M.A., Guerges, A.A., & Ali, S.F. (2009). Genetic signatures of some Egyptian *Clarias gariepinus* populations. *Global Veterinaria*, 3(6): 503-508.
- Skelton, P. (2001). *A Complete Guide to the Freshwater Fishes of Southern Africa*. Struik Publishers, Cape Town. 395 pp.
- Williams, J.K.F., Kubelik, A.R., Livak, K.G., Rafalski, J.A., & Tingey, S.V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, 18:6531-6535. <https://doi:10.1093/nar/18.22.6531>
- Wollebaek, J., Heggenes, J. and Roed, K.H. (2011). Population connectivity: Dam migration mitigations and contemporary site fidelity in Arctic char. *BioMed Central Evolutionary Biology*, 11:207.
- Zheng, X., Cheng, T., Yang, L., Xu, J., Tang, J., Xie, ..., Hu, Z. (2019). Genetic Diversity and DNA Fingerprints of Three Important Aquatic Vegetables by EST-SSR Markers. *Scientific Reports*, 9:14074. <https://doi:10.1038/s41598-019-50569-3>